

# Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

#### A NEW UROMYCES.

BY BYRON D. HALSTED, AMES, IOWA.

The following are the characteristic features of a species of *Uromyces* found near Ames, Iowa, upon leaves of *Leersia Virginica*, Willd.:

UROMYCES DIGITATUS, Halsted.—Sori mostly hypogenous, small, elongated, forming irregular rows; uredospores 22-25x26-30u, elliptical, light yellow, indistinctly echinulate, tips dark; teleutosori shining black, spores wedge-shaped and quite irregular, 17-30x25-33u, pedicel about one-half the length of spore. The broad, free end of the spore is divided into 5-20 sharp or blunt projections, the longer with hyaline tips.

This *Uromyces* approaches *U. acuminatus*, Arth., but differs from it greatly in the wedge or triangular shape of the teleuto-spore and the large number of blunt terminations which cover the broad, free end. These irregular projections make the spore look in shape like a boxing glove; they frequently extend downward along the whole of one side of the spore.

#### NEW LITERATURE.

BY W. A. KELLERMAN.

"Tomato Disease, Cladosporium fulvum." By Chas. B. Plowright. The Gardeners' Chronicle, Oct. 29, 1887.

"THE PREPARATION OF AGARICS FOR THE HERBARIUM." James E. Humphry. Botanical Gazette, November, 1887.

"Beitraegezur Morphologie und Biologie der Uredineen." Von P. Deitel. Botanisches Centralblatt, Band XXII, Nos. 2-5.

"REPORT OF THE MYCOLOGIST, F. L. SCRIBNER." Author's Edition, From the An. Rep. Dept. Agr., 1886.

### ERRATA.

In the November number (pp. 128 and 129) where the word "maculicola" occurs, read "maculiculous."

On p. 126 for "Tricothecium" read "Trichothecium."

## INDEX TO VOLUME III.

#### General Index.

A New Uromyces, B. D. Halsted1	38
Addenda to Index to Genera and Species of Hypocreaceæ	
Additions to Cercospora, Gloeosporium and Cylindrosporium, J. B.	
Flic & R M Evarbart	13